



Superior Livestock Vaccination Programs Designed for Your Management Practices

Management requirements:

- **Please consult your veterinarian or health advisor when selecting the vaccine that qualifies for these programs.**
 - **Always read and follow vaccine label directions.**
 - Always use proper injection sites and techniques. Whenever possible, use Sub Q labeled products.
 - Provide adequate nutrition and minerals to enhance immune system.
 - Include other vaccines or management practices that are unique to your operation or are of value to the calf purchaser.
-

VAC 24

Calves vaccinated on cows at 2 to 4 months of age.

Vaccination requirements:

- 1 dose 7-way, 8-way or 9-way Clostridial.
 - 1 dose viral 5-way (IBR, PI3, BRSV and BVD Type I & II). IBR and PI3 must be chemically altered modified live or modified live with veterinarian's approval. BVD and BRSV can be modified live or killed.
 - 1 dose Mannheimia Haemolytica and/or Pasteurella Multocida.
 - Internal & External Parasite Control Recommended.
-

VAC 34

Calves vaccinated on cows at 2-4 weeks prior to shipping.

Vaccination requirements:

- 2 doses 7-way, 8-way or 9-way Clostridial. One dose at branding and one dose 2-4 weeks prior to shipping.
 - 1 dose viral 5-way (IBR, PI3, BRSV and BVD Type I & II). IBR and PI3 must be chemically altered modified live or modified live with veterinarian's approval. BVD and BRSV can be modified live or killed.
 - 1 dose Mannheimia Haemolytica and/or Pasteurella Multocida.
 - Internal & External Parasite Control Recommended.
-

VAC 34+

Calves vaccinated on cows at branding and 2-4 weeks prior to shipping.

Vaccination requirements:

- 2 doses 7-way, 8-way or 9-way Clostridial. One dose at branding and one dose 2-4 weeks prior to shipping.
 - 2 doses viral 5-way (IBR, PI3, BRSV and BVD Type I & II) at branding and 2-4 weeks prior to shipping. IBR and PI3 must be chemically altered modified live or modified live with veterinarian's approval. BVD and BRSV can be modified live or killed.
 - 1 dose Mannheimia Haemolytica and/or Pasteurella Multocida 2-4 weeks prior to shipping
 - Internal and external parasite control recommended
-

VAC 45

Calves vaccinated twice: 2-4 weeks prior to weaning and at weaning or at weaning and boosted to label instructions. Home Raised and Weaned minimum of 45 days prior to delivery.

Vaccination requirements:

- 2 doses 7-way, 8-way or 9-way Clostridial.
 - 2 doses viral 5-way (IBR, PI3, BRSV and BVD Type I & II). IBR and PI3 must be chemically altered modified live or modified live with veterinarian's approval. BVD and BRSV can be modified live or killed.
 - 1 dose Mannheimia Haemolytica and/or Pasteurella Multocida prior to weaning or at weaning and boosted to label instructions.
 - Internal & External Parasite Control Recommended
-

VAC 45+

Calves vaccinated three times: At branding, prior to or at weaning and boosted to label instructions. Home Raised and Weaned minimum of 45 days prior to delivery.

Vaccination requirements:

- 2 doses 7-way, 8-way or 9-way Clostridial @ branding, prior to or at weaning and boosted to label instructions.
 - 3 doses Modified Live Viral 5 way (IBR, PI3, BRSV, BVD Type I & II) @ branding, prior to weaning or at weaning and boosted to label instructions.
 - 2 doses Mannheimia Haemolytica and/or Pasteurella Multocida @ branding, prior to, or at weaning and boosted to label instructions.
 - Internal & External Parasite Control Required.
-

Vac Precon

Vaccinated twice: At arrival and boosted to label instructions, minimum of 14 days prior to delivery. Purchased cattle weaned minimum of 60 days prior to delivery.

Vaccination requirements:

- 2 doses 7-way, 8-way or 9-way Clostridial on arrival and boosted.
 - 2 doses Modified Live Viral 5 way (IBR, PI3, BRSV, BVD Type I & II) on arrival and boosted.
 - 1 dose Mannheimia Haemolytica and/or Pasteurella Multocida.
 - Internal Parasite Control Recommended, External Parasite Control Recommended.
-